

# Cessnock District Hospital


## 30-day mortality following hospitalisation for seven conditions

The risk-standardised mortality ratio (RSMR) is an indicator that describes, for each hospital's patient cohort, the 'observed' number of deaths divided by the 'expected' number of deaths<sup>1</sup>. The 'expected' number of deaths takes account of the hospital's case mix and is estimated using a statistical model built using the NSW patient population characteristics and outcomes. A ratio of less than 1.0 indicates lower than expected mortality while a ratio greater than 1.0 indicates higher than expected mortality. Small deviations from 1.0 are not considered to be meaningful.


Funnel plots with 95% and 99.8% control limits around the NSW rate are used to identify outlier hospitals – those with 'special cause' variation that may warrant further investigation.

The measure is not designed to enable direct comparisons between hospitals. It assesses each hospital's results given its particular case mix. RSMRs do not distinguish deaths that are avoidable from those that are a reflection of the natural course of illness.


### Risk-standardised mortality ratios (RSMRs) for seven conditions




### RSMRs for three-year periods, seven clinical conditions



### How to interpret the dashboard



### How to interpret a funnel plot




# Cessnock District Hospital

## 30-day mortality following hospitalisation for acute myocardial infarction, July 2012 – June 2015

	This hospital	NSW
Total acute myocardial infarction hospitalisations	105	38,352
Acute myocardial infarction patients		
Presenting patients (index cases) <sup>1</sup>	79	30,488
Patients transferred to another hospital within 30 days	52	14,797
Percentage of patients aged 65+ years* <sup>2</sup>	64.6%	62.3%
Percentage of patients aged 75+ years* <sup>2</sup>	34.2%	38.7%

### Significant patient factors and comorbidities, this hospital, index cases<sup>3</sup>




\*Age was a significant factor in the final model of 30-day mortality following hospitalisation for acute myocardial infarction.

# Cessnock District Hospital

## 30-day mortality following hospitalisation for acute myocardial infarction, July 2012 – June 2015

	This hospital	NSW
Mortality (all causes) among 79 acute myocardial infarction index cases	3 (3.8%)	2,108 (6.9%)
Percentages: index cases who died within 30 days of hospitalisation		
Where deaths occurred:		
Percentage in this hospital	<10 deaths Detailed results not shown	
Percentage in another hospital following transfer		
Percentage after discharge		
When deaths occurred:		
Percentage on day of admission	<10 deaths Detailed results not shown	
Percentage within seven days		


### Cumulative mortality following hospitalisation for acute myocardial infarction, this hospital and NSW<sup>4</sup>



# Cessnock District Hospital

## 30-day mortality following hospitalisation for acute myocardial infarction, July 2012 – June 2015

Acute myocardial infarction risk-standardised mortality ratio by number of expected deaths, NSW public hospitals<sup>5</sup>



### Illustrating the effect of standardisation, July 2012 – June 2015

In order to make fair comparisons, a number of risk adjustments are made to mortality data. These take into account patient factors that influence the likelihood of dying. The table below illustrates the effect of statistical adjustments on this hospital's results.

Unadjusted ratio	Age and sex standardised ratio	Risk-standardised mortality ratio
<b>0.55</b>	<b>0.66</b>	<b>0.97</b>


Ratio: ■ Lower than expected    ■ No different than expected    ■ Higher than expected

The extent to which comorbidities are coded in the patient record may affect risk adjustment. Therefore the 'depth of coding'<sup>6</sup> has been assessed across NSW hospitals. In July 2009 – June 2012, the average depth of coding was 2.6 diagnoses in this hospital and 4.3 in NSW; and in July 2012 – June 2015, there were 2.8 diagnoses in this hospital and 4.8 in NSW.

### Three-yearly RSMR and annual unadjusted mortality rates

The RSMR is calculated on the basis of three years of data. It takes account of differences in patient characteristics so that assessments of hospital performance are fair. To give an indication of results within the three-year period, the figure below shows the RSMR result for July 2012 – June 2015 alongside differences between this hospital and the NSW result for annual unadjusted mortality rates.


RSMR: ○ Lower than expected    ○ No different than expected    ○ Higher than expected



# Cessnock District Hospital

## 30-day mortality following hospitalisation for acute myocardial infarction, July 2012 – June 2015

Acute myocardial infarction, this hospital's risk-standardised mortality ratio, expected mortality rates and observed (unadjusted) mortality rates, July 2000 – June 2015



### Notes

- Data refer to patients who were discharged between July 2012 and June 2015 who were initially admitted to this hospital (regardless of whether they were subsequently transferred) in their last period of care, for an acute and emergency hospitalisation with AMI as principal diagnosis (ICD-10-AM codes I21, excluding I21.9). Deaths are from any cause, in or out of hospital within 30 days of the hospitalisation admission date.
  - Age at admission date.
  - Comorbidities as recorded on patient record, with one year look back from the admission date of the index case. Many are a result of end-organ damage resulting from comorbidities, such as diabetes. The Australian Commission on Safety and Quality in Healthcare comorbidity list was used for acute myocardial infarction, ischaemic stroke, haemorrhagic stroke, pneumonia and hip fracture surgery. The Elixhauser comorbidity list was used for congestive heart failure and chronic obstructive pulmonary disease. STEMI refers to ST-elevation myocardial infarction. Only those conditions that were shown to have a significant impact on mortality ( $P < 0.05$ ) are shown.
  - Cumulative percentage of deaths over the 30 days following admission to hospital for the relevant condition.
  - Results for hospitals with expected deaths  $< 1$  are not shown. Peer hospitals are identified according to the NSW Ministry of Health's peer grouping as of April 2012.
  - The depth of coding has been defined as the average number of secondary diagnosis coded for the index cases. The one year look back method which is used for risk adjustment, to some extent accounts for possible lower depth of coding in some hospitals.
- Details of analyses are available in *Spotlight on Measurement: Measuring 30-day mortality following hospitalisation, 2nd edition*.  
 Data source: SAPHaRI, Centre for Epidemiology and Evidence, NSW Ministry of Health.

# Cessnock District Hospital

**30-day mortality following hospitalisation for ischaemic stroke,  
July 2012 – June 2015**

<50 index hospitalisations,  
results not shown





# Cessnock District Hospital

**30-day mortality following hospitalisation for haemorrhagic stroke,  
July 2012 – June 2015**

<50 index hospitalisations,  
results not shown




# Cessnock District Hospital

## 30-day mortality following hospitalisation for congestive heart failure, July 2012 – June 2015

	This hospital	NSW
Total congestive heart failure hospitalisations	172	40,670
Congestive heart failure patients		
Presenting patients (index cases) <sup>1</sup>	125	27,484
Patients transferred to another hospital within 30 days	13	4,200
Percentage of patients aged 65+ years* <sup>2</sup>	89.6%	90.2%
Percentage of patients aged 75+ years* <sup>2</sup>	70.4%	73.0%

### Significant patient factors and comorbidities, this hospital, index cases<sup>3</sup>




\*Age was a significant factor in the final model of 30-day mortality following hospitalisation for congestive heart failure.

# Cessnock District Hospital

## 30-day mortality following hospitalisation for congestive heart failure, July 2012 – June 2015

	This hospital	NSW
Mortality (all causes) among 125 congestive heart failure index cases	13 (10.4%)	3,793 (13.8%)
Percentages: index cases who died within 30 days of hospitalisation		
Where deaths occurred:		
Percentage in this hospital	53.8%	57.2%
Percentage in another hospital following transfer	0.0%	1.5%
Percentage after discharge	46.2%	41.3%
When deaths occurred:		
Percentage on day of admission	15.4%	5.5%
Percentage within seven days	76.9%	43.9%


### Cumulative mortality following hospitalisation for congestive heart failure, this hospital and NSW<sup>4</sup>



# Cessnock District Hospital

## 30-day mortality following hospitalisation for congestive heart failure, July 2012 – June 2015

Congestive heart failure risk-standardised mortality ratio by number of expected deaths, NSW public hospitals<sup>5</sup>



### Illustrating the effect of standardisation, July 2012 – June 2015

In order to make fair comparisons, a number of risk adjustments are made to mortality data. These take into account patient factors that influence the likelihood of dying. The table below illustrates the effect of statistical adjustments on this hospital's results.

Unadjusted ratio	Age and sex standardised ratio	Risk-standardised mortality ratio
<b>0.75</b>	<b>0.75</b>	<b>0.79</b>

Ratio: ■ Lower than expected    ■ No different than expected    ■ Higher than expected

The extent to which comorbidities are coded in the patient record may affect risk adjustment. Therefore the 'depth of coding'<sup>6</sup> has been assessed across NSW hospitals. In July 2009 – June 2012, the average depth of coding was 3.9 diagnoses in this hospital and 5.1 in NSW; and in July 2012 – June 2015, there were 3.4 diagnoses in this hospital and 6.0 in NSW.

### Three-yearly RSMR and annual unadjusted mortality rates

The RSMR is calculated on the basis of three years of data. It takes account of differences in patient characteristics so that assessments of hospital performance are fair. To give an indication of results within the three-year period, the figure below shows the RSMR result for July 2012 – June 2015 alongside differences between this hospital and the NSW result for annual unadjusted mortality rates.


RSMR: ○ Lower than expected    ○ No different than expected    ○ Higher than expected



# Cessnock District Hospital

## 30-day mortality following hospitalisation for congestive heart failure, July 2012 – June 2015

Congestive heart failure, this hospital's risk-standardised mortality ratio, expected mortality rates and observed (unadjusted) mortality rates, July 2000 – June 2015



### Notes


- Data refer to patients who were discharged between July 2012 and June 2015 who were initially admitted to this hospital (regardless of whether they were subsequently transferred) in their last period of care, for an acute and emergency hospitalisation with congestive heart failure as principal diagnosis (ICD-10-AM codes I11.0, I13.0, I13.2, I50.0, I50.1, I50.9). Deaths are from any cause, in or out of hospital within 30 days of the hospitalisation admission date.
  - Age at admission date.
  - Comorbidities as recorded on patient record, with one year look back from the admission date of the index case. Many are a result of end-organ damage resulting from comorbidities, such as diabetes. The Australian Commission on Safety and Quality in Healthcare comorbidity list was used for acute myocardial infarction, ischaemic stroke, haemorrhagic stroke, pneumonia and hip fracture surgery. The Elixhauser comorbidity list was used for congestive heart failure and chronic obstructive pulmonary disease. STEMI refers to ST-elevation myocardial infarction. Only those conditions that were shown to have a significant impact on mortality ( $P < 0.05$ ) are shown.
  - Cumulative percentage of deaths over the 30 days following admission to hospital for the relevant condition.
  - Results for hospitals with expected deaths  $< 1$  are not shown. Peer hospitals are identified according to the NSW Ministry of Health's peer grouping as of April 2012.
  - The depth of coding has been defined as the average number of secondary diagnosis coded for the index cases. The one year look back method which is used for risk adjustment, to some extent accounts for possible lower depth of coding in some hospitals.
- Details of analyses are available in *Spotlight on Measurement: Measuring 30-day mortality following hospitalisation, 2nd edition*.
- Data source: SAPHaRI, Centre for Epidemiology and Evidence, NSW Ministry of Health.

# Cessnock District Hospital

## 30-day mortality following hospitalisation for pneumonia, July 2012 – June 2015

	This hospital	NSW
Total pneumonia hospitalisations	260	54,478
Pneumonia patients		
Presenting patients (index cases) <sup>1</sup>	230	47,133
Patients transferred to another hospital within 30 days	36	6,564
Percentage of patients aged 65+ years* <sup>2</sup>	65.2%	69.1%
Percentage of patients aged 75+ years* <sup>2</sup>	49.1%	50.0%

### Significant patient factors and comorbidities, this hospital, index cases<sup>3</sup>




\*Age was a significant factor in the final model of 30-day mortality following hospitalisation for pneumonia.

# Cessnock District Hospital

## 30-day mortality following hospitalisation for pneumonia, July 2012 – June 2015

	This hospital	NSW
Mortality (all causes) among 230 pneumonia index cases	27 (11.7%)	5,037 (10.7%)
Percentages: index cases who died within 30 days of hospitalisation		
Where deaths occurred:		
Percentage in this hospital	48.1%	60.3%
Percentage in another hospital following transfer	7.4%	1.4%
Percentage after discharge	44.4%	38.3%
When deaths occurred:		
Percentage on day of admission	7.4%	6.0%
Percentage within seven days	59.3%	51.8%

### Cumulative mortality following hospitalisation for pneumonia, this hospital and NSW<sup>4</sup>






# Cessnock District Hospital

## 30-day mortality following hospitalisation for pneumonia, July 2012 – June 2015

Pneumonia risk-standardised mortality ratio by number of expected deaths, NSW public hospitals<sup>5</sup>



### Illustrating the effect of standardisation, July 2012 – June 2015

In order to make fair comparisons, a number of risk adjustments are made to mortality data. These take into account patient factors that influence the likelihood of dying. The table below illustrates the effect of statistical adjustments on this hospital's results.

Unadjusted ratio	Age and sex standardised ratio	Risk-standardised mortality ratio
<b>1.10</b>	<b>1.21</b>	<b>1.25</b>


Ratio: ■ Lower than expected    ■ No different than expected    ■ Higher than expected

The extent to which comorbidities are coded in the patient record may affect risk adjustment. Therefore the 'depth of coding'<sup>6</sup> has been assessed across NSW hospitals. In July 2009 – June 2012, the average depth of coding was 2.9 diagnoses in this hospital and 3.8 in NSW; and in July 2012 – June 2015, there were 2.8 diagnoses in this hospital and 4.8 in NSW.

### Three-yearly RSMR and annual unadjusted mortality rates

The RSMR is calculated on the basis of three years of data. It takes account of differences in patient characteristics so that assessments of hospital performance are fair. To give an indication of results within the three-year period, the figure below shows the RSMR result for July 2012 – June 2015 alongside differences between this hospital and the NSW result for annual unadjusted mortality rates.


RSMR: ○ Lower than expected    ○ No different than expected    ○ Higher than expected



# Cessnock District Hospital

## 30-day mortality following hospitalisation for pneumonia, July 2012 – June 2015

Pneumonia, this hospital's risk-standardised mortality ratio, expected mortality rates and observed (unadjusted) mortality rates, July 2000 – June 2015



### Notes


- Data refer to patients who were discharged between July 2012 and June 2015 who were initially admitted to this hospital (regardless of whether they were subsequently transferred) in their last period of care, for an acute and emergency hospitalisation with pneumonia as principal diagnosis (ICD-10-AM codes J13, J14, J15, J16, J18). Deaths are from any cause, in or out of hospital within 30 days of the hospitalisation admission date.
  - Age at admission date.
  - Comorbidities as recorded on patient record, with one year look back from the admission date of the index case. Many are a result of end-organ damage resulting from comorbidities, such as diabetes. The Australian Commission on Safety and Quality in Healthcare comorbidity list was used for acute myocardial infarction, ischaemic stroke, haemorrhagic stroke, pneumonia and hip fracture surgery. The Elixhauser comorbidity list was used for congestive heart failure and chronic obstructive pulmonary disease. STEMI refers to ST-elevation myocardial infarction. Only those conditions that were shown to have a significant impact on mortality ( $P < 0.05$ ) are shown.
  - Cumulative percentage of deaths over the 30 days following admission to hospital for the relevant condition.
  - Results for hospitals with expected deaths  $< 1$  are not shown. Peer hospitals are identified according to the NSW Ministry of Health's peer grouping as of April 2012.
  - The depth of coding has been defined as the average number of secondary diagnosis coded for the index cases. The one year look back method which is used for risk adjustment, to some extent accounts for possible lower depth of coding in some hospitals.
- Details of analyses are available in *Spotlight on Measurement: Measuring 30-day mortality following hospitalisation, 2nd edition*.  
 Data source: SAPHaRI, Centre for Epidemiology and Evidence, NSW Ministry of Health.

# Cessnock District Hospital

## 30-day mortality following hospitalisation for chronic obstructive pulmonary disease, July 2012 – June 2015

	This hospital	NSW
Total chronic obstructive pulmonary disease hospitalisations	363	58,675
Chronic obstructive pulmonary disease patients		
Presenting patients (index cases) <sup>1</sup>	192	30,525
Patients transferred to another hospital within 30 days	22	3,337
Percentage of patients aged 65+ years* <sup>2</sup>	78.1%	79.5%
Percentage of patients aged 75+ years* <sup>2</sup>	46.9%	50.7%

### Significant patient factors and comorbidities, this hospital, index cases<sup>3</sup>




\*Age was a significant factor in the final model of 30-day mortality following hospitalisation for chronic obstructive pulmonary disease.

# Cessnock District Hospital

## 30-day mortality following hospitalisation for chronic obstructive pulmonary disease, July 2012 – June 2015

	This hospital	NSW
Mortality (all causes) among 192 chronic obstructive pulmonary disease index cases	20 (10.4%)	3,160 (10.4%)
Percentages: index cases who died within 30 days of hospitalisation		
Where deaths occurred:		
Percentage in this hospital	50.0%	55.8%
Percentage in another hospital following transfer	5.0%	1.4%
Percentage after discharge	45.0%	42.8%
When deaths occurred:		
Percentage on day of admission	10.0%	5.1%
Percentage within seven days	50.0%	47.1%


### Cumulative mortality following hospitalisation for chronic obstructive pulmonary disease, this hospital and NSW<sup>4</sup>



# Cessnock District Hospital

## 30-day mortality following hospitalisation for chronic obstructive pulmonary disease, July 2012 – June 2015

Chronic obstructive pulmonary disease risk-standardised mortality ratio by number of expected deaths, NSW public hospitals<sup>5</sup>



### Illustrating the effect of standardisation, July 2012 – June 2015

In order to make fair comparisons, a number of risk adjustments are made to mortality data. These take into account patient factors that influence the likelihood of dying. The table below illustrates the effect of statistical adjustments on this hospital's results.

Unadjusted ratio	Age and sex standardised ratio	Risk-standardised mortality ratio
<b>1.01</b>	<b>1.06</b>	<b>1.09</b>

Ratio: ■ Lower than expected    ■ No different than expected    ■ Higher than expected

The extent to which comorbidities are coded in the patient record may affect risk adjustment. Therefore the 'depth of coding'<sup>6</sup> has been assessed across NSW hospitals. In July 2009 – June 2012, the average depth of coding was 3.4 diagnoses in this hospital and 3.6 in NSW; and in July 2012 – June 2015, there were 3.0 diagnoses in this hospital and 4.3 in NSW.

### Three-yearly RSMR and annual unadjusted mortality rates

The RSMR is calculated on the basis of three years of data. It takes account of differences in patient characteristics so that assessments of hospital performance are fair. To give an indication of results within the three-year period, the figure below shows the RSMR result for July 2012 – June 2015 alongside differences between this hospital and the NSW result for annual unadjusted mortality rates.


RSMR: ○ Lower than expected    ○ No different than expected    ○ Higher than expected



# Cessnock District Hospital

## 30-day mortality following hospitalisation for chronic obstructive pulmonary disease, July 2012 – June 2015

Chronic obstructive pulmonary disease, this hospital’s risk-standardised mortality ratio, expected mortality rates and observed (unadjusted) mortality rates, July 2000 – June 2015



### Notes

- Data refer to patients who were discharged between July 2012 and June 2015 who were initially admitted to this hospital (regardless of whether they were subsequently transferred) in their last period of care, for an acute and emergency hospitalisation with COPD as principal diagnosis (ICD-10-AM code J41, J42, J43, J44, J47, and J20 and J40 if accompanied by J41, J42, J43, J44 and J47 in any secondary diagnoses). Deaths are from any cause, in or out of hospital within 30 days of the hospitalisation admission date.
  - Age at admission date.
  - Comorbidities as recorded on patient record, with one year look back from the admission date of the index case. Many are a result of end-organ damage resulting from comorbidities, such as diabetes. The Australian Commission on Safety and Quality in Healthcare comorbidity list was used for acute myocardial infarction, ischaemic stroke, haemorrhagic stroke, pneumonia and hip fracture surgery. The Elixhauser comorbidity list was used for congestive heart failure and chronic obstructive pulmonary disease. STEMI refers to ST-elevation myocardial infarction. Only those conditions that were shown to have a significant impact on mortality ( $P < 0.05$ ) are shown.
  - Cumulative percentage of deaths over the 30 days following admission to hospital for the relevant condition.
  - Results for hospitals with expected deaths  $< 1$  are not shown. Peer hospitals are identified according to the NSW Ministry of Health’s peer grouping as of April 2012.
  - The depth of coding has been defined as the average number of secondary diagnosis coded for the index cases. The one year look back method which is used for risk adjustment, to some extent accounts for possible lower depth of coding in some hospitals.
- Details of analyses are available in *Spotlight on Measurement: Measuring 30-day mortality following hospitalisation, 2nd edition*.  
 Data source: SAPHaRI, Centre for Epidemiology and Evidence, NSW Ministry of Health.

# Cessnock District Hospital

**30-day mortality following hospitalisation for hip fracture surgery,  
July 2012 – June 2015**

<50 index hospitalisations,  
results not shown

