# Bateman's Bay District Hospital: Activity profiles, potentially avoidable admissions for COPD and CHF


July 2009 to June 2010

	last year	one year ago
5,176	4,715	10%
2,291	2,227	3%
592	571	4%
78	88	-11%
61	65	-6%
	2,291 592 78	5,176 4,715   2,291 2,227   592 571   78 88

### Rates per 1,000 select medical hospitalisations<sup>5</sup>

COPD <sup>4</sup>	This period (2009-10)	Last period (2008-09)	Peer group (2009-10)	CHF <sup>4</sup>	This period (2009-10)	Last period (2008-09)	Peer group (2009-10)
Actual	34.1	39.7	41.4	Actual	26.7	29.3	22.5
Standardised	13.7	*	38.8	Standardised	*	*	24.1
95% CI	(6.1 - 22.7)	*	(36.4 - 41.3)	95% CI	*	*	(21.8 - 26.4)

# Bateman's Bay District Hospital: Patient profiles, potentially avoidable admissions for COPD and CHF July 2009 to June 2010




Same period Change since


## Bateman's Bay District Hospital: Hospital stay profiles, potentially avoidable admissions for COPD and CHF July 2009 to June 2010

COPD admissions	4 This period (2009-10)	Last period (2008-09)	Peer group (2009-10)	CHF admissions <sup>4</sup>	This period (2009-10)	Last period (2008-09)	Peer group (2009-10)	
% Unplanned <sup>10</sup>	100%	100%	97%	% Unplanned <sup>10</sup>	100%	100%	97%	
% from ED <sup>‡</sup>	82%	93%	77%	% from ED <sup>‡</sup>	82%	80%	70%	
COPD average length of stay in days <sup>4</sup>				CHF average lengt	th of stay in day	ys <sup>4</sup>		
Actual	4.1	5.2	5.8	Actual	5.6	4.6	6.7	

#### Length of stay profiles



## Bateman's Bay District Hospital: Total bed days, potentially avoidable admissions for COPD and CHF July 2005 to June 2010 (monthly)



- Suppressed: relative standard error ≥ 40%.
- (†) Interpret with caution: 30% ≤ relative standard error < 40%.
- A hospitalisation corresponds to one 'episode of care'. A single patient can have multiple hospitalisations in the time period.
- 2. Excludes hospitalisations for renal dialysis, obstetrics, psychiatry, and all surgery.
- 3. Potentially preventable hospitalisations as defined in: The health of the people of NSW Report of the Chief Health Officer 2010.
- 4. For criteria used for COPD / CHF admissions, see Technical Supplement at www.bhi.nsw.gov.au
- 5. Rates are directly standardised to the NSW admitted patient population 2009-10 on the basis of age, sex, and socioeconomic status.
- 6. Index of Relative Social Disadvantage (IRSD), as defined by the Australian Bureau of Statistics.
- Smoking status as recorded on admission: termed current if patient smoked tobacco within last month; previous if patient smoked tobacco in the past but excluding last month.
- We use the term Aboriginal, rather than Aboriginal and Torres Strait Islander in line with NSW Health usage, which recognises that Aboriginal people are the original inhabitants of NSW.
- Postcode of usual residence at time of admission classified as outer regional or remote. 9.
- 10. Admissions with emergency status of 'non-emergency / planned'.

(\*) Emergency Department

Note: Only records with valid and non-missing data are included in each analysis.

Note: Length of stay and total bed days calculated on 'acute' episodes only (comprising 98.7% COPD and 97.4% CHF PAAs).

Note: Additional detail and definitions can be found in the Bureau's Technical Supplement at www.bhi.nsw.gov.au

Source: Admitted Patient Data Collection. Health Information Exchange, NSW Health. Data extracted on 28 February 2011.